

49. In the Claims:

Please cancel claims 1-17.

Please add the following claims:

- 10001733.110201
- a3
- 7
18. (New) A tablet comprising
- (a) from about 3000 to about 9000 FCC lactase units and
- (b) from about 25 to about 70 weight percent microcrystalline cellulose.
- 8
19. (New) A tablet according to claim 18 further comprising a member selected from the group consisting of dextrose, mannitol, a lubricant, cornstarch, calcium phosphate, sodium citrate, calcium sulfate, calcium stearate, stearic acid, glyceryl monostearate, glyceryl distearate, sorbitol, gelatin, a gum, and mixtures thereof.
- 9
20. (New) A tablet according to claim 19 comprising dextrose, sodium citrate, and a lubricant.
- 10
21. (New) A tablet according to claim 19 comprising manitol, dextrose, sodium citrate, and a gum.
- 11
22. (New) A tablet according to claim 21 wherein the gum is selected from the group consisting of carboxymethyl cellulose, methyl cellulose, alginate, dextran, acacia gum, karaya gum, locust bean gum, and tragacanth.
- 12
23. (New) A tablet according to claim 22 wherein the gum is carboxymethyl cellulose.
- 13
24. (New) A tablet according to claim 19 wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.
- 14
25. (New) A tablet according to claim 20 wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.
- 15
26. (New) A tablet according to claim 24 wherein the lubricant is magnesium stearate.
- 16
27. (New) A tablet according to claim 25 wherein the lubricant is magnesium stearate.

¹⁷
~~28.~~ (New) A tablet according to claim ⁴~~19~~ wherein the lubricant is present in an amount of from about 0.25 to about 6 % by weight.

¹⁸
~~29.~~ (New) A tablet according to claim ⁹~~20~~ wherein the lubricant is present in an amount of from about 0.25 to about 6 % by weight.

¹⁹
~~30.~~ (New) A tablet comprising

- (a) about 3000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 25 to about 70 weight percent mannitol;
- (d) from about 0.25 to about 6 weight percent magnesium stearate;
- (a) carboxymethyl cellulose;
- (b) dextrose; and
- (c) sodium citrate.

²⁰
~~31.~~ (New) A tablet comprising

- (a) about 9000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 25 to about 70 weight percent mannitol;
- (d) from about 0.25 to about 6 weight percent magnesium stearate;
- (e) carboxymethyl cellulose;
- (f) dextrose; and
- (g) sodium citrate.

²¹
~~32.~~ (New) A tablet comprising

- (a) about 9000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 0.25 to about 6 weight percent magnesium stearate;
- (d) dextrose; and
- (e) sodium citrate.

²²
~~33.~~ (New) A composition comprising

- (a) from about 3000 to about 9000 FCC lactase units and
- (b) from about 25 to about 70 weight percent microcrystalline cellulose.

10001733 110201

a3
Cont

²³
~~34~~

(New) A composition according to claim ²²~~33~~ further comprising a member selected from the group consisting of dextrose, mannitol, a lubricant, cornstarch, calcium phosphate, sodium citrate, calcium sulfate, calcium stearate, stearic acid, glyceryl monostearate, glyceryl distearate, sorbitol, gelatin, a gum, and mixtures thereof.

²⁴
~~35~~

(New) A composition according to claim ²³~~34~~ comprising dextrose, sodium citrate, and a lubricant.

²⁵
~~36~~

(New) A composition according to claim ²³~~34~~ comprising mannitol, dextrose, sodium citrate, and a gum.

²⁶
~~37~~

(New) A composition according to claim ²⁵~~36~~ wherein the gum is selected from the group consisting of carboxymethyl cellulose, methyl cellulose, alginate, dextran, acacia gum, karaya gum, locust bean gum, and tragacanth.

²⁷
~~38~~

(New) A composition according to claim ²⁶~~37~~ wherein the gum is carboxymethyl cellulose.

²⁸
~~39~~

(New) A composition according to claim ²³~~34~~ wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.

²⁹
~~40~~

(New) A composition according to claim ²⁴~~35~~ wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.

³⁰
~~41~~

(New) A composition according to claim ²⁹~~40~~ wherein the lubricant is magnesium stearate.

³¹
~~42~~

(New) A composition according to claim ²⁴~~35~~ wherein the lubricant is magnesium stearate.

³²
~~43~~

(New) A composition according to claim ²³~~34~~ wherein the lubricant is present in an amount of from about 0.25 to about 5 % by weight.

³³
~~44~~

(New) A composition according to claim ²⁴~~35~~ wherein the lubricant is present in an amount of from about 0.25 to about 5 % by weight.

10001733-110201

a3
cont.

34
45.

(New) A composition comprising

- (a) about 3000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 25 to about 70 weight percent mannitol;
- (d) from about 0.25 to about 6 weight percent magnesium stearate;
- (e) carboxymethyl cellulose;
- (f) dextrose; and
- (g) sodium citrate.

35
46.

(New) A composition comprising

- (a) about 9000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 25 to about 70 weight percent mannitol;
- (d) from about 0.25 to about 6 weight percent magnesium stearate;
- (e) carboxymethyl cellulose;
- (f) dextrose; and
- (g) sodium citrate.

36
47.

(New) A composition comprising

- (a) about 9000 FCC lactase units;
- (b) from about 25 to about 70 weight percent microcrystalline cellulose;
- (c) from about 0.25 to about 6 weight percent magnesium stearate;
- (d) dextrose; and
- (e) sodium citrate.

37
48.

(New) A process for making a tablet comprising the following steps in order

- (a) mixing lactase and microcrystalline cellulose to form a preblend;
- (b) adding to the preblend a member selected from the group consisting of dextrose, mannitol, a lubricant, cornstarch, calcium phosphate, sodium citrate, calcium sulfate, calcium stearate, stearic acid, glyceryl monostearate, glyceryl distearate, sorbitol, gelatin, a gum, and mixtures thereof to form a mixture; and
- (c) compressing the mixture to form a tablet.

38
49.

(New) A process according to claim 48 wherein dextrose, sodium citrate, and a lubricant are added to the preblend.

39
50.

(New) A process according to claim 48 wherein manitol, dextrose, sodium citrate, and a gum are added to the preblend.

40
51.

(New) A process according to claim 33 wherein the gum is selected from the group consisting of carboxymethyl cellulose, methyl cellulose, alginate, dextran, acacia gum, karaya gum, locust bean gum, and tragacanth.

41
52.

(New) A process according to claim 34 wherein the gum is carboxymethyl cellulose.

42
53.

(New) A process according to claim 31 wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.

43
54.

(New) A process according to claim 32 wherein the lubricant is selected from the group consisting of magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oils, and mixtures thereof.

44
55.

(New) A process according to claim 36 wherein the lubricant is magnesium stearate.

45
56.

(New) A process according to claim 37 wherein the lubricant is magnesium stearate.

46
57.

(New) A process according to claim 36 wherein the lubricant is present in an amount of from about 0.25 to about 6 % by weight.

47
58.

(New) A process according to claim 37 wherein the lubricant is present in an amount of from about 0.25 to about 6 % by weight.

48
59.

(New) A process for making a tablet comprising

- (a) mixing an amount lactase having a potency of between about 14,000 and about 100,000 FCC units and about 25 to about 85 weight percent microcrystalline cellulose to form a preblend;
- (b) adding from about 25 to about 70 weight percent of the tablet of mannitol; from about 0.25 to about 5 weight percent of the tablet of magnesium stearate; carboxymethyl cellulose; dextrose; and sodium citrate to form a mixture;

10001733-110201

a3
Cont

(c) and compressing the mixture to form the tablet.

49
~~60~~

(New) A process for making a tablet comprising

(a) mixing an amount lactase having a potency of between about 14,000 and about 100,000 FCC units and about 25 to about 85 weight percent

microcrystalline cellulose to form a preblend;

(b) adding from about 0.25 to about 5 weight percent of the tablet of magnesium stearate; dextrose; and sodium citrate to form a mixture; and

(c) compressing the mixture to form the tablet.

10001733-110201